HAB MONITORING REPORT

From: 7/5/2016 To: 7/5/2016

Fish and Wildlife Research Institute

Collecting agency: EBAP

Collected by: Staff

Sample condition: Preserved

| HAB ID | Location | County | Lat/Lon | | Depth | Temp | Sal | DO | pН | Species | cells/liter |
|---|--|--------|---------------------|-------|-------|-------|-------|--------|------|-----------------------|-------------|
| Original ID | | | (DD.dddd) | | (m) | (C) | (ppt) | (mg/L) | | | |
| Sample Date | | | | | | | | | | | |
| HABW160706-031 FDEP EBV001 7/5/2016 | Matanzas Pass (Estero Bay) | Lee | 26.4577 -81.9532 | 07:22 | 0.5 | 32.00 | 24.30 | 4.54 | 7.95 | | |
| | schen, K. on 7/11/2016 | | | | | | | | | Karenia brevis | 0 |
| Comments: wind | | | | | | | | | | Pseudo-nitzschia spp. | 0 |
| | | | | | | | | | | Pyrodinium bahamense | 0 |
| HABW160706-032 FDEP EBV003 7/5/2016 | Estero River; mouth of (Estero Bay) | Lee | 26.4294 -81.8580 | 06:48 | 0.5 | 30.60 | 25.67 | 2.02 | 8.29 | | |
| | schen, K. on 7/11/2016 | | | | | | | | | Karenia brevis | 0 |
| Comments: winds SE@2-3mph | | | | | | | | | | Pseudo-nitzschia spp. | 0 |
| | | | | | | | | | | Pyrodinium bahamense | 0 |
| HABW160706-033 FDEP EBV004 7/5/2016 | Carl Johnson Park Boat Ramp (Estero Bay) | Lee | 26.3936 -81.8655 | 06:52 | 0.5 | 30.50 | 30.69 | 2.72 | 8.00 | | |
| | schen, K. on 7/11/2016 | | | | | | | | | Karenia brevis | 0 |
| Comments: wind | Is NE@4mph | | | | | | | | | Pseudo-nitzschia spp. | 0 |
| | | | | | | | | | | Pyrodinium bahamense | 0 |
| HABW160706-034 FDEP EBV005 7/5/2016 | Pelican Bay Nature Park Pier (Estero Bay) | Lee | 26.3584 -81.8375 | 06:40 | 0.5 | 29.90 | 28.71 | 2.61 | 7.94 | , | |
| | schen, K. on 7/11/2016 | | | | | | | | | Karenia brevis | 0 |
| Comments: winds E@2-3mph | | | | | | | | | | Pseudo-nitzschia spp. | 0 |
| | | | | | | | | | | Pyrodinium bahamense | 0 |
| HABW160706-035 FDEP EBV006 7/5/2016 | Coon Key; N of (Estero Bay) | Lee | 26.4287 -81.8832 | 06:46 | 0.5 | 30.00 | 26.26 | 2.76 | 8.15 | | |
| Analyzed by: Hens | schen, K. on 7/11/2016 | | | | | | | | | Karenia brevis | 0 |
| Comments: wind | ls NE@4-7mph | | | | | | | | | Pseudo-nitzschia spp. | 0 |
| | | | | | | | | | | Pyrodinium bahamense | 0 |

| HAB ID | Location | County | Lat/Lon | Time | Depth | Temp | Sal | DO | pН | Species | cells/liter |
|---|-------------------------------|--------|---------------------|-------|-------|-------|-------|--------|------|-----------------------|-------------|
| Original ID | | | (DD.dddd) |) | (m) | (C) | (ppt) | (mg/L) | | | |
| Sample Date | | | | | | | | | | | |
| HABW160706-036 FDEP EBV007 7/5/2016 | Mound House Dock (Estero Bay) | Lee | 26.4462 -81.9272 | 07:48 | 0.5 | 32.20 | 24.24 | 5.14 | 7.87 | | |
| Analyzed by: Hen | schen, K. on 7/11/2016 | | | | | | | | | Karenia brevis | 0 |
| Comments: winds E@2-3mph | | | | | | | | | | Pseudo-nitzschia spp. | 0 |
| | | | | | | | | | | Pyrodinium bahamense | 0 |
| HABW160706-037 FDEP EBERS2 7/5/2016 | Estero River; upstream | Lee | 26.4386 -81.8400 | 07:20 | 0.5 | 29.20 | 0.97 | 3.28 | 7.40 | | |
| | schen, K. on 7/11/2016 | | | | | | | | | Karenia brevis | 0 |
| Comments: seas E@0-1mph | | | | | | | | | | Pseudo-nitzschia spp. | 0 |
| | | | | | | | | | | Pyrodinium bahamense | 0 |

HAB MONITORING REPORT

From: 7/5/2016 To: 7/5/2016

Fish and Wildlife Research Institute



Collecting agency: FDEP

Collected by: Staff

Sample condition: Preserved

| HAB ID | Location | County | Lat/Lon Time | Depth | Temp | Sal | DO | рН | Species | cells/liter |
|-------------------------------|---------------------------------------|--------|---------------------|-------|------|-------|--------|----|-----------------------|-------------|
| Original ID | | | (DD.dddd) | (m) | (C) | (ppt) | (mg/L) |) | | |
| Sample Date | | | | | | | | | | |
| HABW160706-038 FDEP CHV009 | Burnt Store Marina (Charlotte Harbor) | Lee | 26.7614 -82.0611 | 0.5 | | | | | | |
| 7/5/2016 | | | | | | | | | | |
| | schen, K. on 7/11/2016 | | | | | | | | Karenia brevis | 0 |
| Comments: | | | | | | | | | Pseudo-nitzschia spp. | 0 |
| | | | | | | | | | Pyrodinium bahamense | 0 |

| Description | Karenia brevis abundance | Possible effects (<i>Karenia brevis</i> only) |
|-----------------------------|-------------------------------|---|
| NOT PRESENT - BACKGROUND | 0 - 1,000 cells/L | no effects anticipated |
| VERY LOW | > 1,000 - 10,000 cells/L | possible respiratory irritation; shellfish harvesting closures ≥ 5,000 cells/L |
| LOW | > 10,000 - 100,000 cells/L | respiratory irritation; possible fish kills; probable detection of surface chlorophyll by satellites at upper range of cell abundance |
| MEDIUM | > 100,000 - 1,000,000 cells/L | respiratory irritation; probable fish kills; detection of surface chlorophyll by satellites |
| нідн | > 1,000,000 cells/L | as above, plus water discoloration |

The above report is distributed by the Harmful Algal Bloom (HAB) Group at the Fish and Wildlife Research Institute of the Florida Fish and Wildlife Conservation Commission. The report is intended to (1) provide timely information on HABs in Florida waters to partner agencies and (2) facilitate communication among individuals who direct response activities to address public health concerns. We report on the abundance of *Karenia brevis*, *Pyrodinium bahamense* and *Pseudonitzschia* species. *Karenia brevis*, the Florida red tide organism, produces neurotoxins called brevetoxins that can kill fish and other marine life. Brevetoxins may cause respiratory irritation in beachgoers and Neurotoxic Shellfish Poisoning in humans that consume contaminated shellfish. *Pyrodinium bahamense* produces saxitoxins that can cause Paralytic Shellfish Poisoning or Saxitoxin Puffer Fish Poisoning in humans if contaminated shellfish or puffer fish are consumed. Some, but not all, species of *Pseudo-nitzschia* produce domoic acid, which can cause Amnesic Shellfish Poisoning in humans if contaminated shellfish are consumed. Blooms of *Pseudo-nitzschia* spp. (≥ 1,000,000 cells/L) frequently occur in Florida's marine and estuarine waters. For information on red tide related human health issues, please refer to the <u>Department of Health Aquatic Toxins Program</u>.

State-wide status reports of Karenia brevis abundance including interactive Google Maps are provided weekly by our group. Shellfish harvesting area status maps are provided by the Division of Aquaculture. Gulf Coast beach conditions can be found at Mote Marine Laboratory's Beach Conditions Report. A full list of red tide related hotlines and information sources can be found here. Data for other species can be requested at any time by sending an inquiry to HABData@MyFWC.com. To learn more about HAB monitoring and research in Florida, please visit MyFWC.com/RedTide and Facebook.com/FLHABs.

DISCLAIMER: While every practical step has been taken to provide accurate information in these reports, the need for rapid distribution precludes extensive review. Further, reports are generated with limited interpretation and do not necessarily reflect all scientific observations.



Matanzas Pass (Estero Bay)

Mound House Dock (Estero Bay)

Estero River; upstream

Coon Key; N of (Estero Bay)

Estero River; mouth of (Estero Bay)

Carl Johnson Park Boat Ramp (Estero Bay)

Karenia brevis (cells/liter)

not present/background (0-1,000)
very low (>1,000-10,000)
low (>10,000-100,000)
medium (>100,000-1,000,000)
high (>1,000,000)

Pelican Bay Nature Park Pier (Estero Bay)


Lee

Google earth

Image © 2016 TerraMetrics

4.23 km





Google earth

Karenia brevis (cells/liter)

not present/background (0-1,000)
very low (>1,000-10,000)
low (>10,000-100,000)
medium (>100,000-1,000,000)
high (>1,000,000)

2527 m