

HAB MONITORING REPORT
From: 4/3/2017 To: 4/3/2017
Fish and Wildlife Research Institute



HAB ID Original ID Sample Date	Location	County	Lat/Lon (DD.ddddd)	Time	Depth (m)	Temp (C)	Sal (ppt)	DO (mg/L)	pH	Species	cells/liter
HABW170404-014 FDEP EBV003 4/3/2017	Estero River; mouth of (Estero Bay)	Lee	26.4294 -81.8580	07:22	0.5	25.90	35.51	3.30	7.51		
Collected by: Franklin, N. of EBAP; Preserved Analyzed by: Markley, L. on 4/4/2017 Comments: S wind at 2-3mph; ripples; partly cloudy; incoming tide										<i>Karenia brevis</i> <i>Pseudo-nitzschia spp.</i> <i>Pyrodinium bahamense</i>	0 1,333 0
HABW170404-015 FDEP EBV004 4/3/2017	Carl Johnson Park Boat Ramp (Estero Bay)	Lee	26.3936 -81.8655	08:13	0.5	25.90	35.65	4.11	7.99		
Collected by: Winter, T. of EBAP; Preserved Analyzed by: KellerAbbe, S. on 4/4/2017 Comments: E wind at 4-7mph; waves; partly cloudy; high slack										<i>Karenia brevis</i> <i>Pseudo-nitzschia spp.</i> <i>Pyrodinium bahamense</i>	9,000 264,633 0
HABW170404-016 FDEP EBV005 4/3/2017	Pelican Bay Nature Park Pier (Estero Bay)	Lee	26.3584 -81.8375	07:22	0.5	25.70	34.14	3.66	7.66		
Collected by: Sims, C. of EBAP; Preserved Analyzed by: KellerAbbe, S. on 4/4/2017 Comments: SE wind at 4-7mph; ripples; fog/haze; incoming tide										<i>Karenia brevis</i> <i>Pseudo-nitzschia spp.</i> <i>Pyrodinium bahamense</i>	333 17,333 0
HABW170404-017 FDEP EBV006 4/3/2017	Coon Key; N of (Estero Bay)	Lee	26.4287 -81.8832	07:02	0.5	26.20	34.88	5.26	7.30		
Collected by: Volunteer(s) of EBAP; Preserved Analyzed by: KellerAbbe, S. on 4/4/2017 Comments: SE wind at 4-7mph; ripples; overcast; high slack										<i>Karenia brevis</i> <i>Pseudo-nitzschia spp.</i> <i>Pyrodinium bahamense</i>	3,333 272,417 0

NOTE: Blank field = not measured

HAB ID	Location	County	Lat/Lon (DD.dddd)	Time	Depth (m)	Temp (C)	Sal (ppt)	DO (mg/L)	pH	Species	cells/liter
HABW170404-018	Mound House Dock (Estero Bay)	Lee	26.4462 -81.9272	07:15	0.5	26.00	35.09	4.26	7.93		
FDEP EBV007											
4/3/2017											
	Collected by: Flynn, R. of EBAP; Preserved									<i>Karenia brevis</i>	0
	Analyzed by: Markley, L. on 4/4/2017									<i>Pseudo-nitzschia spp.</i>	21,667
	Comments: SE wind at 8-12mph; waves; partly cloudy; incoming tide									<i>Pyrodinium bahamense</i>	0
HABW170404-019	Estero River; upstream	Lee	26.4386 -81.8400	07:40	0.5	26.50	29.34	2.47	7.14		
FDEP EBERS2											
4/3/2017											
	Collected by: Fretwell of EBAP; Preserved									<i>Karenia brevis</i>	0
	Analyzed by: Henschen, K. on 4/4/2017									<i>Pseudo-nitzschia spp.</i>	0
	Comments: SE wind at 4-7mph; ripples; overcast; incoming tide									<i>Pyrodinium bahamense</i>	0

NOTE: Blank field = not measured

Description	<i>Karenia brevis</i> abundance	Possible effects (<i>Karenia brevis</i> only)
NOT PRESENT - BACKGROUND	0 - 1,000 cells/L	no effects anticipated
VERY LOW	> 1,000 - 10,000 cells/L	possible respiratory irritation; shellfish harvesting closures \geq 5,000 cells/L
LOW	> 10,000 - 100,000 cells/L	respiratory irritation; possible fish kills; probable detection of surface chlorophyll by satellites at upper range of cell abundance
MEDIUM	> 100,000 - 1,000,000 cells/L	respiratory irritation; probable fish kills; detection of surface chlorophyll by satellites
HIGH	> 1,000,000 cells/L	as above, plus water discoloration

The above report is distributed by the Harmful Algal Bloom (HAB) Group at the Fish and Wildlife Research Institute of the Florida Fish and Wildlife Conservation Commission. The report is intended to (1) provide timely information on HABs in Florida waters to partner agencies and (2) facilitate communication among individuals who direct response activities to address public health concerns. We report on the abundance of [Karenia brevis](#), [Pyrodinium bahamense](#) and [Pseudo-nitzschia](#) species. *Karenia brevis*, the Florida red tide organism, produces neurotoxins called brevetoxins that can kill fish and other marine life. Brevetoxins may cause respiratory irritation in beachgoers and Neurotoxic Shellfish Poisoning in humans that consume contaminated shellfish. *Pyrodinium bahamense* produces saxitoxins that can cause Paralytic Shellfish Poisoning or Saxitoxin Puffer Fish Poisoning in humans if contaminated shellfish or puffer fish are consumed. Some, but not all, species of *Pseudo-nitzschia* produce domoic acid, which can cause Amnesic Shellfish Poisoning in humans if contaminated shellfish are consumed. Blooms of *Pseudo-nitzschia* spp. (\geq 1,000,000 cells/L) frequently occur in Florida's marine and estuarine waters. For information on red tide related human health issues, please refer to the [Department of Health Aquatic Toxins Program](#).

[State-wide status reports](#) of *Karenia brevis* abundance including interactive Google Maps are provided weekly by our group. [Shellfish harvesting area status maps](#) are provided by the Division of Aquaculture. Gulf Coast beach conditions can be found at [Mote Marine Laboratory's Beach Conditions Report](#). A full list of red tide related hotlines and information sources can be found [here](#). Data for other species can be requested at any time by sending an inquiry to HABData@MyFWC.com. To learn more about HAB monitoring and research in Florida, please visit MyFWC.com/RedTide and Facebook.com/FLHABs.

DISCLAIMER: While every practical step has been taken to provide accurate information in these reports, the need for rapid distribution precludes extensive review. Further, reports are generated with limited interpretation and do not necessarily reflect all scientific observations.



Lee

Mound House Dock (Estero Bay)

Estero River; upstream

Coon Key; N of (Estero Bay)

Estero River; mouth of (Estero Bay)

Carl Johnson Park Boat Ramp (Estero Bay)

Pelican Bay Nature Park Pier (Estero Bay)

***Karenia brevis* (cells/liter)**

- not present/background (0-1,000)
- very low (>1,000-10,000)
- low (>10,000-100,000)
- medium (>100,000-1,000,000)
- high (>1,000,000)

5.65 km

Image © 2016 TerraMetrics
Data SIO, NOAA, U.S. Navy, NGA, GEBCO

Google earth