HAB MONITORING REPORT

From: 10/7/2019 To: 10/7/2019

Collected by: Volunteer(s)
Collecting agency: EBAP
Sample condition: Preserved

Fish and Wildlife Research Institute



HAB ID Original ID Sample Date	Location	County	Lat/Lon (DD.dddd)	Time)	Depth (m)	Temp (C)	Sal (ppt)	DO (mg/L)	pН	Species	cells/lite
HABW191009-024 FDEP EBV001 10/7/2019	Matanzas Pass (Estero Bay)	Lee	26.4577 -81.9532	08:04	0.5	27.90	33.67	1.06	7.66		
	KellerAbbe, S. on 10/9/2019 Wind E @ 8 - 12 mph, overcast, air temp 26.5 C, secch m, tide high slack, water color yellow-green	ni 19.5								Karenia brevis Pseudo-nitzschia spp. Pyrodinium bahamense	23,000 9,667 0
HABW191009-025 FDEP EBV004 10/7/2019	Carl Johnson Park Boat Ramp (Estero Bay)	Lee	26.3936 -81.8655	07:39	0.5	27.60	34.78	3.70	7.88		
	Henschen, K. on 10/9/2019 Wind S @ 4-7 mph, overcast, air temp 23.7 C,, tide incoming, secchi = 1.4 m, water color yellow-green									Karenia brevis Pseudo-nitzschia spp. Pyrodinium bahamense	35,333 14,000 0
HABW191009-026 FDEP EBV005 10/7/2019	Pelican Bay Nature Park Pier (Estero Bay)	Lee	26.3584 -81.8375	07:25	0.5	28.70	32.82	4.01	7.78		
	Henschen, K. on 10/9/2019 Wind NE @ 4 - 7 mph, overcast, air temp 25.3 C, tide incoming, secchi = 0.95 m, water color yellow-green									Karenia brevis Pseudo-nitzschia spp. Pyrodinium bahamense	667 13,667 0
HABW191009-027 FDEP EBV006 10/7/2019	Coon Key; N of (Estero Bay)	Lee	26.4287 -81.8832	07:19	0.5	27.90	33.67	4.58	7.83		
	Henschen, K. on 10/9/2019 Wind SE @ 8 - 12 mph, overcast, air temp 24.6 C, tide slack, secchi = 0.95 m, water color green-brown	high								Karenia brevis Pseudo-nitzschia spp. Pyrodinium bahamense	333 0 0
HABW191009-028 FDEP EBV007 10/7/2019	Mound House Dock (Estero Bay)	Lee	26.4462 -81.9272	07:47	0.5	28.00	32.96	3.98	7.70	,	
Analyzed by:	Henschen, K. on 10/9/2019 Wind NE @ 4 - 7 mph, overcast, air temp 23.8 C, tide incoming, secchi = 1.8 m, water color yellow-brown									Karenia brevis Pseudo-nitzschia spp. Pyrodinium bahamense	333 10,667 0

HAB ID	Location	County	Lat/Lon	Time	Depth	Temp	Sal	DO (11)	рН	Species	cells/liter
Original ID			(DD.dddd)	(m)	(C)	(ppt)	(mg/L))		
Sample Date											
HABW191009-029 FDEP EBERS2 10/7/2019	Estero River; upstream	Lee	26.4386 -81.8400	07:50	0.5	27.80	25.18	3.45	7.21		
Comments:	Henschen, K. on 10/9/2019 Wind E @ 2 - 3 mph, overcast, air temp 24.5 C, tide outgoing, secchi = 1.7 m, water color green-brown									Karenia brevis Pseudo-nitzschia spp. Pyrodinium bahamense	0 0 0

NOTE: Blank field = not measured

Description	Karenia brevis abundance	Possible effects (<i>Karenia brevis</i> only)
NOT PRESENT - BACKGROUND	0 - 1,000 cells/L	no effects anticipated
VERY LOW	> 1,000 - 10,000 cells/L	possible respiratory irritation; shellfish harvesting closures ≥ 5,000 cells/L
LOW	> 10,000 - 100,000 cells/L	respiratory irritation; possible fish kills; probable detection of surface chlorophyll by satellites at upper range of cell abundance
MEDIUM	> 100,000 - 1,000,000 cells/L	respiratory irritation; probable fish kills; detection of surface chlorophyll by satellites
HIGH	> 1,000,000 cells/L	as above, plus water discoloration

The above report is distributed by the Harmful Algal Bloom (HAB) Group at the Fish and Wildlife Research Institute of the Florida Fish and Wildlife Conservation Commission. The report is intended to (1) provide timely information on HABs in Florida waters to partner agencies and (2) facilitate communication among individuals who direct response activities to address public health concerns. We report on the abundance of *Karenia brevis*, *Pyrodinium bahamense* and *Pseudonitzschia* species. *Karenia brevis*, the Florida red tide organism, produces neurotoxins called brevetoxins that can kill fish and other marine life. Brevetoxins may cause respiratory irritation in beachgoers and Neurotoxic Shellfish Poisoning in humans that consume contaminated shellfish. *Pyrodinium bahamense* produces saxitoxins that can cause Paralytic Shellfish Poisoning or Saxitoxin Puffer Fish Poisoning in humans if contaminated shellfish or puffer fish are consumed. Some, but not all, species of *Pseudo-nitzschia* produce domoic acid, which can cause Amnesic Shellfish Poisoning in humans if contaminated shellfish are consumed. Blooms of *Pseudo-nitzschia* spp. (≥ 1,000,000 cells/L) frequently occur in Florida's marine and estuarine waters. For information on red tide related human health issues, please refer to the Department of Health Aquatic Toxins Program.

State-wide status reports of *Karenia brevis* abundance including interactive Google Maps are provided weekly by our group. Shellfish harvesting area status maps are provided by the Division of Aquaculture. Gulf Coast beach conditions can be found at Mote Marine Laboratory's Beach Conditions Report. A full list of red tide related hotlines and information sources can be found here. Data for other species can be requested at any time by sending an inquiry to HABData@MyFWC.com/Research/redtide and Facebook.com/FLHABs.

DISCLAIMER: While every practical step has been taken to provide accurate information in these reports, the need for rapid distribution precludes extensive review. Further, reports are generated with limited interpretation and do not necessarily reflect all scientific observations.

