HAB MONITORING REPORT

From: 7/6/2020 To: 7/6/2020

Fish and Wildlife Research Institute

Collected by: Volunteer(s) Collecting agency: EBAP

Sample condition: Preserved

HAB ID	Location	County	Lat/Lon	Time	Depth	Temp	Sal	DO	pН	Species	cells/liter
Original ID			(DD.dddd)		(m)	(C)	(ppt)	(mg/L)			
Sample Date											
HABW200707-035 FDEP EBV004 7/6/2020	Carl Johnson Park Boat Ramp (Estero Bay)	Lee	26.3936 -81.8655		0.5	30.50	33.32	2.33	8.61		
Analyzed by:	Thurmond, R. on 7/8/2020									Karenia brevis	0
Comments:	Wind SW @ 4-7 mph, drizzle, air 24.6 C, tide low slack									Pseudo-nitzschia spp.	6,667
	secchi 1.0, green-brown; heavy rain prior to sampling, speed increased after.	, wind								Pyrodinium bahamense	0
HABW200707-048 FDEP EBV006 7/6/2020	Coon Key; N of (Estero Bay)	Lee	26.4287 -81.8832	06:41	0.5	31.00	32.62	4.83	8.10		
Analyzed by:	Thurmond, R. on 7/8/2020									Karenia brevis	0
Comments:	Wind S @ 3.9 mph, partly cloudy, air temp 25.7 C, tid									Pseudo-nitzschia spp.	5,333
	incoming, secchi = 1.2 m, water color medium brown									Pyrodinium bahamense	0
HABW200707-049 FDEP EBV007 7/6/2020	Mound House Dock (Estero Bay)	Lee	26.4462 -81.9272	06:12	0.5	31.80	32.49	4.50	8.18		
	Thurmond, R. on 7/8/2020									Karenia brevis	0
Comments:	Wind S @ 8 - 12 mph, partly cloudy, air 28.4 C, incom	ling								Pseudo-nitzschia spp.	1,667
	tide, sechhi = 1.6 m, water green-brown									Pyrodinium bahamense	0

HAB MONITORING REPORT

From: 7/6/2020 To: 7/6/2020

Fish and Wildlife Research Institute

ON THE REAL PROPERTY OF THE RE

Sample condition: Preserved

Collected by: Kowitch, L. Collecting agency: PC

HAB ID	Location	County	Lat/Lon	Time	Depth	Temp	Sal	DO	рН	Species	cells/liter
Original ID			(DD.dddd	I.	(m)	(C)	(ppt)	(mg/L)			
Sample Date											
HABW200707-052	Big Carlos Pass	Lee	26.4056 -81.8833	06:50	0.5	26.00					
7/6/2020											
Analyzed by: Thu	rmond, R. on 7/8/2020									Karenia brevis	0
Comments: Heavy rain prior to sampling										Pseudo-nitzschia spp.	7,667
										Pyrodinium bahamense	0
HABW200707-053	Hogue Channel	Lee	26.3579 -81.8563	09:00	0.5	27.00					
7/6/2020											
Analyzed by: Thu	rmond, R. on 7/8/2020									Karenia brevis	0
Comments: Hear	vy rain prior to sampling									Pseudo-nitzschia spp.	5,333
										Pyrodinium bahamense	0

Description	Karenia brevis abundance	Possible effects (<i>Karenia brevis</i> only)
NOT PRESENT - BACKGROUND	0 - 1,000 cells/L	no effects anticipated
VERY LOW	> 1,000 - 10,000 cells/L	possible respiratory irritation; shellfish harvesting closures \geq 5,000 cells/L
LOW	> 10,000 - 100,000 cells/L	respiratory irritation; possible fish kills; probable detection of surface chlorophyll by satellites at upper range of cell abundance
MEDIUM	> 100,000 - 1,000,000 cells/L	respiratory irritation; probable fish kills; detection of surface chlorophyll by satellites
нідн	> 1,000,000 cells/L	as above, plus water discoloration

The above report is distributed by the Harmful Algal Bloom (HAB) Group at the Fish and Wildlife Research Institute of the Florida Fish and Wildlife Conservation Commission. The report is intended to (1) provide timely information on HABs in Florida waters to partner agencies and (2) facilitate communication among individuals who direct response activities to address public health concerns. We report on the abundance of *Karenia brevis*, *Pyrodinium bahamense* and *Pseudonitzschia* species. <u>Karenia brevis</u>, the Florida red tide organism, produces neurotoxins called brevetoxins that can kill fish and other marine life. Brevetoxins may cause respiratory irritation in beachgoers and Neurotoxic Shellfish Poisoning in humans that consume contaminated shellfish. <u>Pyrodinium bahamense</u> produces saxitoxins that can cause Paralytic Shellfish Poisoning or Saxitoxin Puffer Fish Poisoning in humans if contaminated shellfish or puffer fish are consumed. Some, but not all, species of <u>Pseudo-nitzschia</u> produce domoic acid, which can cause Amnesic Shellfish Poisoning in humans if contaminated shellfish are consumed. Blooms of *Pseudo-nitzschia* spp. (≥ 1,000,000 cells/L) frequently occur in Florida's marine and estuarine waters. For information on red tide related human health issues, please refer to the <u>Department of Health Aquatic Toxins Program</u>.

State-wide status reports of Karenia brevis abundance including interactive Google Maps are provided weekly by our group. Shellfish harvesting area status maps are provided by the Division of Aquaculture. Gulf Coast beach conditions can be found at Mote Marine Laboratory's Beach Conditions Report. A full list of red tide related hotlines and information sources can be found here. Data for other species can be requested at any time by sending an inquiry to HABData@MyFWC.com. To learn more about HAB monitoring and research in Florida, please visit MyFWC.com/Research/redtide and Facebook.com/FLHABs.

DISCLAIMER: While every practical step has been taken to provide accurate information in these reports, the need for rapid distribution precludes extensive review. Further, reports are generated with limited interpretation and do not necessarily reflect all scientific observations.



Mound House Dock

Coon Key; N of

Big Carlos Pass

Carl Johnson Park Boat Ramp

Karenia brevis (cells/liter)

not present/background (0-1,000)
very low (>1,000-10,000)
low (>10,000-100,000)
medium (>100,000-1,000,000)

high (>1,000,000)

Google⁻earth

Image © 2020 TerraMetrics

Hogue Channel

3 mi