HAB MONITORING REPORT

From: 6/6/2022 To: 6/6/2022

Fish and Wildlife Research Institute



Collecting agency: FDEP-EBAP **Sample condition:** Preserved

Collected by: Volunteer(s)

HAB ID Original ID Sample Date	Location	County	Lat/Lon (DD.dddd		Depth (m)	Temp (C)	Sal (ppt)	DO (mg/L)	рН	Species	cells/liter
HABW220607-037 FDEP EBV003 6/6/2022	Estero River; mouth of (Estero Bay)	Lee	26.4294 -81.8580	06:39	0.5	28.90	21.47	4.90	7.83		
	Markley, L. on 6/7/2022 Wind SW @ 0-1 mph, partly cloudy, air 24.7 C; incom tide, secchi = 1.00, water green brown	ning								Karenia brevis Pseudo-nitzschia spp. Pyrodinium bahamense	0 11,000 0
HABW220607-038 FDEP EBV004 6/6/2022	Carl Johnson Park Boat Ramp (Estero Bay)	Lee	26.3936 -81.8655	06:45	0.5	28.80	28.16	3.17	7.95		
	Markley, L. on $6/7/2022$ Wind NE @ 2-3 mph, partly cloudy, air 24.6 C; incom secchi = 1.3 m, water yellow green	ing tide,								Karenia brevis Pseudo-nitzschia spp. Pyrodinium bahamense	0 112,000 0
HABW220607-039 FDEP EBV005 6/6/2022	Pelican Bay Nature Park Pier (Estero Bay)	Lee	26.3584 -81.8375	06:53	0.5	28.80	26.37	4.41	7.81		
	Henschen, K. on $6/7/2022$ Wind S @ 0-1 mph, partly cloudy skies, air 24.3 C; tio incoming, secchi = 1.20, water green brown	de								Karenia brevis Pseudo-nitzschia spp. Pyrodinium bahamense	0 1,333 0
HABW220607-040 FDEP EBV007 6/6/2022	Mound House Dock (Estero Bay)	Lee	26.4462 -81.9272	06:15	0.5	28.50	21.28	4.56	7.84		
	Henschen, K. on $6/7/2022$ Wind NE @ 0-1 mph, partly cloudy skies, air 24.7 C, to incoming, secchi = 1.65 m, water yellow brown	tide								Karenia brevis Pseudo-nitzschia spp. Pyrodinium bahamense	0 39,667 0
HABW220607-041 FDEP EBERS2 6/6/2022	Estero River; upstream	Lee	26.4386 -81.8400	07:20	0.5	27.10	0.75	2.65	7.23		
Analyzed by:	Henschen, K. on 6/7/2022 No wind, partly cloudy skies, air 24.0 C; incoming tide = 1.6 m, water dark brown	e, secchi								Karenia brevis Pseudo-nitzschia spp. Pyrodinium bahamense	0 0 0

HAB ID	Location	County	Lat/Lon		Depth	Temp	Sal	DO	pН	Species	cells/liter
Original ID			(DD.dddd)	(m)	(C)	(ppt)	(mg/L)			
Sample Date											
HABW220607-042 FDEP EBV001 6/6/2022	Matanzas Pass (Estero Bay)	Lee	26.4577 -81.9532	06:30	0.5	28.40	27.26	4.33	7.99		
	Henschen, K. on 6/7/2022 Wind E @ 2-3 mph, partly cloudy skies, air 24.7; tide incoming, secchi = 1.6 m, water green brown									Karenia brevis Pseudo-nitzschia spp. Pyrodinium bahamense	0 101,667 0

Description	Karenia brevis abundance	Possible effects (<i>Karenia brevis</i> only)
NOT PRESENT - BACKGROUND	0 - 1,000 cells/L	no effects anticipated
VERY LOW	> 1,000 - 10,000 cells/L	possible respiratory irritation; shellfish harvesting closures ≥ 5,000 cells/L
LOW	> 10,000 - 100,000 cells/L	respiratory irritation; possible fish kills; probable detection of surface chlorophyll by satellites at upper range of cell abundance
MEDIUM	> 100,000 - 1,000,000 cells/L	respiratory irritation; probable fish kills; detection of surface chlorophyll by satellites
нідн	> 1,000,000 cells/L	as above, plus water discoloration

The above report is distributed by the Harmful Algal Bloom (HAB) Group at the Fish and Wildlife Research Institute of the Florida Fish and Wildlife Conservation Commission. The report is intended to (1) provide timely information on HABs in Florida waters to partner agencies and (2) facilitate communication among individuals who direct response activities to address public health concerns. We report on the abundance of *Karenia brevis*, *Pyrodinium bahamense* and *Pseudonitzschia* species. *Karenia brevis*, the Florida red tide organism, produces neurotoxins called brevetoxins that can kill fish and other marine life. Brevetoxins may cause respiratory irritation in beachgoers and Neurotoxic Shellfish Poisoning in humans that consume contaminated shellfish. *Pyrodinium bahamense* produces saxitoxins that can cause Paralytic Shellfish Poisoning or Saxitoxin Puffer Fish Poisoning in humans if contaminated shellfish or puffer fish are consumed. Some, but not all, species of *Pseudo-nitzschia* produce domoic acid, which can cause Amnesic Shellfish Poisoning in humans if contaminated shellfish are consumed. Blooms of *Pseudo-nitzschia* spp. (≥ 1,000,000 cells/L) frequently occur in Florida's marine and estuarine waters. For information on red tide related human health issues, please refer to the <u>Department of Health Aquatic Toxins Program</u>.

State-wide status reports of Karenia brevis abundance including interactive Google Maps are provided weekly by our group. Shellfish harvesting area status maps are provided by the Division of Aquaculture. Gulf Coast beach conditions can be found at Mote Marine Laboratory's Beach Conditions Report. A full list of red tide related hotlines and information sources can be found here. Data for other species can be requested at any time by sending an inquiry to HABData@MyFWC.com. To learn more about HAB monitoring and research in Florida, please visit MyFWC.com/Research/redtide and Facebook.com/FLHABs.

DISCLAIMER: While every practical step has been taken to provide accurate information in these reports, the need for rapid distribution precludes extensive review. Further, reports are generated with limited interpretation and do not necessarily reflect all scientific observations.

CONSET DATION COMMIT

Matanzas Pass

Mound House Dock

Estero River; upstream

Estero River; mouth of

Carl Johnson Park Boat Ramp

Karenia brevis (cells/liter)

not present/background (0-1,000)
very low (>1,000-10,000)
low (>10,000-100,000)
medium (>100,000-1,000,000)

high (>1,000,000)

Pelican Bay Nature Park Pier

Google Earth

.ee

Data SIO, NOAA, U.S. Navy, NGA, GEBCO Image © 2022 TerraMetrics